

INFANT FORMULA

PRODUCT SPECIFICATION



Product description

This product is TRUVITA Infant Formula developed to meet the nutritional requirement of infants aged 0–6 months. This product is formulated with minerals, vitamins and other essential nutrients that support infant's growth and development

Complies with GB10765 (National food safety standard, Infant Formula) and GB 14880 (National food safety standard, Standard for using of nutritional fortification substances in foods)

Claimable country of origin: Made in New Zealand

Product Classification: Infant Formula

Appearance

Pale yellow Powder, No different flavour and taste.

Ingredients

Lactose, Vegetable Oils, Skim Milk, Whey Powder, **Minerals** (Calcium Carbonate, Calcium Phosphate Dibasic, Potassium Chloride, Dipotassium Hydrogen Phosphate, Sodium Chloride, Magnesium Chloride, Ferrous Sulphate, Zinc Sulphate, Copper Sulphate, Manganese Sulphate, Potassium Iodide, Sodium Selenite), Galacto-Oligosaccharides (GOS), Arachidonic Acid (AA, from *Mortierella alpina*) Powder, Docosahexaenoic Acid (DHA, from *Schizochytrium* sp.) Powder, **Vitamins** (L-Ascorbic Acid (Vitamin C), dl- α -Tocopherol Acetate (Vitamin E), Vitamin A Acetate, Cholecalciferol (Vitamin D₃), Niacinamide, Thiamine Hydrochloride (Vitamin B₁), Calcium Pantothenate (Vitamin B₅), Phytonadione (Vitamin K₁), Pyridoxine Hydrochloride (Vitamin B₆), Folic Acid, Cyanocobalamin (Vitamin B₁₂), D-Biotin, Riboflavin (Vitamin B₂), β -Carotene), Lactoferrin, Choline Chloride, Taurine, L-Carnitine, **Nucleotides** (Cytidine 5'-monophosphate, Uridine 5'-monophosphate disodium, Adenosine 5'-monophosphate, Inosine 5'-monophosphate disodium, Guanosine 5'-monophosphate disodium), Inositol.

Nutrition Information

Nutrient composition	Unit	Average Quantity per		
		100 g	100 mL	100 kJ
Energy	kJ	2120	286	100
Protein	g	12	1.6	0.57
Whey : Casein		60 : 40		
Lactoferrin	mg	90	12	4.2
Fat	g	26	3.5	1.2
Linoleic Acid (LA)	mg	4860	656	229
α -Linolenic Acid (ALA)	mg	413	56	19
LA/ALA Ratio		12 : 1		
ARA	mg	60	8.1	2.8
DHA	mg	30	4.1	1.4
Carbohydrate	g	55	7.4	2.6
Lactose	g	54	7.3	2.5
GOS	mg	600	81	28
Vitamins				
Vitamin A	μ g	411	55	19
Vitamin B ₁	μ g	446	60	21
Vitamin B ₂	μ g	604	82	29
Niacin	μ g	2890	390	136
Pantothenic Acid	μ g	3410	460	161
Vitamin B ₆	μ g	315	42	15
Vitamin B ₁₂	μ g	2.0	0.27	0.09
Vitamin C	mg	79	11	3.7
Vitamin D	μ g	7.6	1.0	0.36
Vitamin E	mg	10.0	1.4	0.47
Vitamin K ₁	μ g	37	5.0	1.7
Folic Acid	μ g	104	14	4.9
Biotin	μ g	23	3.1	1.1

INFANT FORMULA

PRODUCT SPECIFICATION



Nutrient composition	Unit	Average Quantity per		
		100g	100 mL	100 kJ
Minerals				
Calcium	mg	485	65	23
Chloride	mg	438	59	21
Copper	µg	311	42	15
Iodine	µg	100	14	4.7
Iron	mg	5.0	0.68	0.24
Magnesium	mg	50	6.8	2.4
Manganese	µg	131	18	6.2
Phosphorus	mg	309	42	14
Potassium	mg	587	79	28
Selenium	µg	17	2.3	0.80
Sodium	mg	186	25	8.8
Zinc	mg	4.0	0.54	0.19
Ca/P Ratio		1.6 : 1		
Nucleotides	mg	21.7	2.93	1.02
Others				
L-Carnitine	mg	13	1.8	0.61
Taurine	mg	28	3.8	1.3
Choline	mg	66	8.9	3.1
Inositol	mg	30	4.0	1.4

Direction for Use (How to Prepare)

- 1) Wash your hands thoroughly with soap and clean warm water before preparing the formula.
- 2) Boil in clean water for 5 minutes – the milk bottles, teat, cap and any utensils or containers required.
- 3) Boil clean drinking water for 5 minutes, and cool until it is lukewarm.
- 4) Pour the exact amount of lukewarm water into the milk bottle as detailed in the 'Feeding Guide'
- 5) Use only the enclosed scoop, fill it with the levelled and measured amount of milk powder. Add to the water.
- 6) Shake the mixture well and test the temperature of the milk with a few drops on the inside of your wrist before feeding. Make sure prepared formula is warm, not hot. Discard any formula left after a feed.
- 7) If multiple bottles of formula are prepared, each bottle should be prepared separately. Bottles must be refrigerated and consumed within 24 hours. Prepared formula left over after this time must be discarded.
- 8) To prevent scalding or burning do not microwave this products.

Feeding Guide

Infant's age	Cooled boiled water (mL)	Level scoops of formula*	Feeds per Day
Up to 2 weeks	80	2	6 – 7
2 weeks – 2 months	120	3	5 – 6
2 – 6 months	200	5	4 – 5

* 1 level scoop = 6.2 g

Note: 13.5 g powder added to 87 mL yields approximately 100 mL.

INFANT FORMULA

PRODUCT SPECIFICATION



Important Notice

Breast milk is best for babies. Before you decide to use this product, consult your doctor or health worker for advice.

This product may be used for infants from birth to 6 months when mothers breast milk is insufficient or unavailable. For correct feeding of your infant, follow the instruction of infant and child nutritionists, or refer to the feeding guide to match formula intake with your infant's appetite.

Warning

Follow instructions exactly. Prepare bottles and teats as directed. Do not change proportions of powder except on medical advice. Incorrect preparation can make your baby very ill.

Storage and Shelf Life

Store TRUVITA Infant Formula at room temperature in dry conditions. Product can be kept for 2 years in room temperature when it is not opened. After opening, seal the lid well to maintain freshness. Consume this product within 4 weeks after opening. For use by date, date of manufacture, batch number, and manufacturing facility, refer to the can base.